



Red Edge as a Potential Index for Detecting Differences in Plant Nitrogen Status in Winter Wheat

Journal:	<i>Journal of Plant Nutrition</i>
Manuscript ID:	LPLA-2010-0106.R1
Manuscript Type:	Original Articles
Date Submitted by the Author:	27-Oct-2010
Complete List of Authors:	Kanke, Yumiko; Oklahoma State University, Plant and Soil Sciences Raun, William; Oklahoma State University, Plant and soil Sciences Solie, John; Oklahoma State University, Biosystems and Agricultural Engineering Stone, Marvin; Oklahoma State University, Biosystems and Agricultural Engineering Taylor, Randal; Oklahoma State University, Biosystems and Agricultural Engineering
Keywords:	Wheat < Crops, Nitrogen < Macronutrients, Soil Fertility, sensors, NDVI, red edge

SCHOLARONE™
Manuscripts

INTRODUCTION

Nitrogen (N) is one of the major limiting mineral nutrients for plant growth. Raun and Johnson (1999) estimated worldwide cereal NUE to be approximately 33%. Due to the continuous increase in fertilizer costs and growing environmental concerns associated with fertilizer use, application of N fertilizer according to plant need has become increasingly popular due to its potential for increasing NUE and reducing input costs. To determine the optimum N rate based on plant need, optical sensing technologies have been developed to detect N status in plants. Normalized difference vegetative index (NDVI) computed from optical sensing is one of the most widely used vegetative indices for the evaluation of plant N status.

There are however, some drawbacks to using NDVI. It has been reported to have low sensitivity at high chlorophyll content or abundant biomass. Gitelson et al. (2002) listed several possible reasons for low sensitivity of NDVI. Decreased NIR reflectance was associated with changes in leaf orientation from one growth stage to the next, reduction in chlorophyll content at senescence, and increasing soil moisture. This also results in the poor estimation of biomass once soil is covered by vegetation (Clevers and Jongschaap, 2001).

To overcome these limitations, wavebands called “red-edge” were employed as new spectra to evaluate plant N conditions. Red-edge wavebands are between RED (670nm) and NIR (780nm). These bands were shown to have greater sensitivity at higher chlorophyll content, which was detected as greener biomass. REP is influenced by chlorophyll content, leaf mesophyll structure, and LAI (Meer and Jong, 2006). On the other hand, leaf orientation, solar angle and soil background had a small influence on REP. Also by combining plant growth models with REP, they improved the estimation of yield in sugarcane (Meer and Jong, 2006).

Several methods have been developed to find the REP. One is detecting the inflection point of the reflectance curve which is the maximum slope between RED and NIR (Meer and Jong, 2006). It uses the first derivative analysis to detect REP (Chen and Elvidge, 1993). Another method is the linear method

1
2
3 which estimates the maximum inflection point by interpolating among four bands; 670, 700, 740, and 780
4 nm (Guyot and Baret, 1988). Shafri et al (2006) reported that this linear method has more soil background
5 noise than the Lagrangian interpolation technique which is based on the spectrum derivative analysis.
6
7 With the linear method, the overestimation of REP by about 10 nm was found compared with the first
8 derivative method (Dawson and Curran, 1998). However, Dawson and Curran (1998) also reported that
9 both methods were correlated at differing chlorophyll levels and the correlation coefficient of REP
10 determined by different methods was high ($R^2 > 0.99$). Mutanga and Skidmore (2007) drew attention to
11 the double points for the red-edge especially in the high N treated plant using the first derivative method.
12 If double REP exists, the linear method is not appropriate to detect the red-edge position. Cho and
13 Skidmore (2006) also developed another method where REP is determined by the intersection of the far-
14 red and red lines on the first derivative reflectance. REP determined by this method increased the linear
15 relationship with N concentration compared with the first derivative method or linear method.
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 Oklahoma State University (OSU) has been developing algorithms for N fertilization for various
31 crops since the early 1990's. The algorithms are based on the use of an optical, active light, handheld
32 GreenSeeker™ sensor which detects the fraction of light being reflected from the plant. The OSU
33 algorithm uses GreenSeeker™ NDVI values as the key input for calculation of the optimum mid-season
34 N fertilization rate. Due to reported limitations indicating NDVI is insensitive to high chlorophyll
35 concentrations or plant biomass, it was necessary to evaluate the potential of red edge for detecting N
36 differences. From an agronomic perspective, it is crucial that timing of N application as well as the rate of
37 N fertilizer be considered. The optimum time to make a decision for mid-season N application in winter
38 wheat is at Feekes 4 to 5 (Large, 1954). It was reported that when mid-season N was applied between
39 Feekes 3 and 4, there was no yield loss (Boman et al., 1995). After Feekes 4, tissue damage and lower
40 forage yields were detected from having applied foliar N. Rapid N uptake occurs between Feekes 2 to 4
41 and by Feekes 7, wheat takes up more than a third of the total accumulated (Waldren and Flowerday,
42 1979). Therefore, in winter wheat, it is essential to determine the N rate for mid-season application at or
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 before Feekes 4. At this time, the wheat plant does not completely cover the ground; there it can be
4
5 assumed that NDVI is still sensitive to plant biomass. It is, thus, essential to investigate how NDVI and
6
7 REP behave differently for early season growth of winter wheat.
8
9

10
11 The SPAD meter is also commercially available and has been used for detecting N differences in
12
13 winter wheat (Fox R.H. et al., 1994). This device emits light at 650 and 940 nm. The transmittance ratio is
14
15 then used for estimation of chlorophyll content which is ultimately related with the N status in plants.
16
17 Therefore, it is essential to evaluate how it behaves differently from REP.
18
19

20
21 The objective of this paper was to determine whether the red-edge index has the potential to be a
22
23 useful index for detecting differences in N status for winter wheat compared to NDVI and SPAD meter.
24
25
26
27

28 **MATERIALS AND METHODS**

29
30 A spectrometer and a chlorophyll meter (SPAD-502) were used to collect data in winter wheat.
31
32 Measurements were taken at different growth stages (Feekes growth stages 4, 5, 7, and 10) for two
33
34 cropping seasons.
35
36

37
38 Data were collected from long-term winter wheat experimental plots located at Stillwater
39
40 (Experiment # 222) and Perkins (N & P Study), Oklahoma. Experiment # 222 was established in 1969
41
42 under conventional tillage on a Kirkland silt loam (fine, mixed, superactive, thermic Udertic Paleustoll).
43
44 The N & P study was initiated in 1996, also under conventional tillage on a Teller sandy loam (fine-
45
46 loamy, mixed, thermic Udic Argiustoll). These experiments are long-term N-P-K trials consisting of
47
48 thirteen treatments (Experiment # 222) with four replications, and twelve treatments (N & P study) with
49
50 three replications, respectively. Both were arranged in a randomized complete block design (RCBD).
51
52 Four N treatments (0, 40, 90, and 135 N kg/ha) and (0, 56, 112 and 168 kg N/ha) were evaluated in
53
54 Experiment # 222 and the N & P study, respectively.
55
56
57
58
59
60

Two instruments were used to obtain data for this study: the Minolta SPAD 502 meter and an Ocean Optics USB4000 spectrometer. All of the readings were taken from a 1 m² area in each plot. The Minolta SPAD 502 chlorophyll meter determines the relative amount of chlorophyll by measuring light transmitted or absorbed by plant leaves. The SPAD 502 is a compact meter that measures chlorophyll using optical density differences at two wavelengths (650 nm and 940 nm) with a measurement area of 2 mm x 3 mm. Twenty SPAD readings were randomly taken from winter wheat plant leaves within the 1 m² sampling area, and subsequently averaged. The Ocean Optics USB4000 spectrometer operates with Spectrasuite (cross-platform spectroscopy software) to measure reflectance. This spectrometer can detect reflectance from 200-1100 nm at a high resolution (optical resolution of 1.5 nm full width half maximum). Reflectance of the plant canopy was computed by (the reflected light from the surface of the plant canopy minus black measurement to eliminate noise)/(incident light minus black measurement). Incident light was determined by measuring reflectance of a 1m² white plate composed of barium sulfate. Dark current was measured by covering the sensor with a cap and fabric material.

Spectral Calculation

Red-edge position

For the REP, two methods were applied: Derivative method by curve fitting techniques and the linear method. For the derivative method, spectrometer reflectance from 650 nm to 750 nm were collected and transported into SYSTAT Table Curve 2.D. software and interpolated using a curve fitting formula . By using the formula, the maximum point of the first derivative was recorded as REP. For the linear method (Figure 1), the interpretation by Clevers (1994) was used.

$$\rho_{REP} = \frac{\rho_{670} + \rho_{780}}{2} \quad (1)$$

$$REP = 700 + 40 * \frac{(\rho_{REP} - \rho_{700})}{(\rho_{740} - \rho_{700})} \quad (2)$$

NDVI and Simple ratio

NDVI and simple ratio computed as follows.

$$NDVI_{RED} = \frac{\rho_{780} - \rho_{670}}{\rho_{780} + \rho_{670}} \quad (3)$$

$$NDVI_{GREEN} = \frac{\rho_{780} - \rho_{560}}{\rho_{780} + \rho_{560}} \quad (4)$$

$$SR_{RED} = \frac{\rho_{780}}{\rho_{670}} \quad (5)$$

$$SR_{GREEN} = \frac{\rho_{780}}{\rho_{560}} \quad (6)$$

RESULTS

Stillwater

In 2007, spectrometer measurements at Feekes 10 were excluded due to measurement errors.

Overall, REP was highly correlated with all others at all growth stages (Table 1). High correlation between REP and $NDVI_{RED}$ was found. In both 2008 and 2009, the relationship between REP and SPAD increased with advancing plant growth. Compared with $NDVI_{GREEN}$, $NDVI_{RED}$ had a better relationship with REP. REP tended to have a higher relationship with SR_{RED} and SR_{GREEN} (simple ratio) than $NDVI_{RED}$ and $NDVI_{GREEN}$ (normalized index). It was found that there was slightly higher correlation between REP and SPAD compared with correlation between $NDVI_{RED}$ and SPAD (Tables 1 and 2). It possibly indicates that REP has higher sensitivity to chlorophyll concentration than $NDVI_{RED}$. SR_{RED} is widely recognized to detect plant biomass on the ground (Bellairs et al., 1996, Serrano et al., 2000 and

1
2
3 Barbar et al., 2006). $NDVI_{RED}$ tended to have higher correlation with SR_{RED} than REP which indicated
4
5 that $NDVI_{RED}$ could be more sensitive to plant biomass.
6
7

8
9 There was a significant influence of N rate on REP at all growth stages both in 2008 and 2009
10 crop years ($p < 0.05$, Figure 2). Nitrogen rate and REP had significant linear relationships at all growth
11 stages ($p < 0.05$). Quadratic relationships between N rates and REP were found at Feekes 7 in 2008 and
12 Feekes 4 and 5 in 2009 ($p < 0.05$). The REP shifted to longer wavelengths with the increase in N rates. The
13 range of REP increased as plant growth progressed in 2008 (4.33, 3.88 and 5.39 nm at Feekes 4, 5, and 7
14 respectively). At high N rates, shifts of REP to longer wavelengths were clearly detected as growth stage
15 increased.
16
17
18
19
20
21
22

23
24 There was a significant influence of N rate on $NDVI_{RED}$. Nitrogen rate and $NDVI_{RED}$ had a
25 significant linear relationship at all growth stages. Quadratic relationships between N rates and $NDVI_{RED}$
26 were not found in 2008 but found at Feekes 4 and 5 in 2009 ($\alpha = 0.05$). In 2009, Experiment #222 had a
27 heavy rye grass infestation, and as a result, plant growth was limited. This could explain the low $NDVI_{RED}$
28 values in 2009 compared with 2008 at whole growth stages. $NDVI_{RED}$ tended to be more sensitive for
29 detecting differences in growth stages than REP (Figure 2). These results showed that $NDVI_{RED}$ better
30 detected differences in plant growth at the same N rates, especially at early growth stages.
31
32
33
34
35
36
37
38
39
40
41
42

43 Perkins

44
45 At Feekes 7 in 2008, spectrometer data were excluded due to measurement error. At Feekes 5 in 2009, all
46 data were excluded due to measurement error.
47
48
49

50
51 Overall, REP was highly correlated with all others at all growth stages except SPAD in 2008
52 (Table 3). REP and $NDVI_{RED}$ were highly correlated. As recorded in Stillwater the correlation between
53 REP and SPAD increased with advancing plant growth. REP tended to be more highly correlated with
54 SR_{RED} and SR_{GREEN} (simple ratio) than $NDVI_{RED}$ and $NDVI_{GREEN}$ (normalized index) at Stillwater but this
55
56
57
58
59
60

1
2
3 was not detected at Perkins. Higher correlation between REP and SPAD was found compared with
4
5 correlation between NDVI and SPAD at both Stillwater and Perkins (Tables 1 and 2, for Stillwater, and
6
7 Tables 3 and 4 for Perkins), suggesting that REP has higher sensitivity to chlorophyll concentration than
8
9 NDVI.
10

11
12
13 There was a significant influence of N rate on REP only at Feekes 10 in 2008 and at all growth
14
15 stages in 2009 ($p < 0.05$, Figure 3). Nitrogen rate and REP were linearly correlated at all growth stages
16
17 ($p < 0.05$). The REP shifted to longer wavelengths with increased N rates but the shifts were not found
18
19 with advancing plant growth. The range of REP increased as plant growth progressed in 2008 (2.10, 3.0.3
20
21 and 5.64 nm at Feekes 4, 5, and 10 respectively) and in 2009 (3.12, 8.41, and 7.76 at Feekes 4, 7 and 10
22
23 respectively).
24

25
26
27 There was a significant influence of N rate on $NDVI_{RED}$ at all growth stages and in both years
28
29 excluding Feekes 4 in 2008 ($p < 0.05$)(Figure 3). A quadratic relationship between N rate and $NDVI_{RED}$
30
31 was found at Feekes 4 and 7 in 2009 ($p < 0.05$). In general sensor values for $NDVI_{RED}$ detected differences
32
33 in plant N response and growth stage for both years. Once plant growth was sufficient to provide
34
35 complete plant cover (beyond Feekes 7), it was still possible to detect N rate differences using $NDVI_{RED}$
36
37 or REP (Figure 3, Feekes 10).
38
39
40
41

42 DISCUSSION

43
44 First, the relationship between NDVI and REP needs to be discussed. Researchers have noted
45
46 that REP is an alternative index and could have higher sensitivity under dense green biomass. In this
47
48 research, NDVI and REP were linearly related. The relationship between NDVI and N rate, and REP and
49
50 N rate were processed in Table Curve 2D v5.01.01 to acquire linear, logarithmic, exponential, and
51
52 sigmoid models. Then parameters of each model including, r^2 , adjusted r^2 , standard error, and F score,
53
54 were tested with a t-test comparison to determine if REP and NDVI were different in terms of detecting
55
56 differences in N rates. Results in Table 5 and 6 show that there was no significant difference between
57
58
59
60

1
2
3 indices for the parameters reported, excluding standard error for all models. The significant difference
4
5 between indices for the standard error could be explained by the difference in units between REP and
6
7 NDVI. REP is described by a 3 digit number (e.g., 724 nm) while NDVI is described by 3 decimals (e.g.,
8
9 0.455). Therefore, the standard deviation of REP should be larger and as a result, significant differences
10
11 between indices were recorded for the standard error. Other than that, there were no significant
12
13 differences of indices between r^2 , adjusted r^2 , and F score. There was also no difference between NDVI
14
15 and REP models for detecting differences in plant N.
16
17

18
19 In Oklahoma, farmers apply N, mid season at Feekes 4 and 5. But, in general, they need to make
20
21 the fertilizer N rate decision before Feekes 5. At that time, the ground is not fully covered by biomass
22
23 (Figure 4). Therefore, low sensitivity of NDVI to plant biomass is not a problem. However with the
24
25 decreased biomass on the ground, other problems arise. With a decrease in plant cover on the ground, the
26
27 soil is more exposed and it might increase the noise for NDVI. In Figure 5, the relationship between REP
28
29 and NDVI is described at Feekes 4. At the high N rate where more surface biomass was expected, the
30
31 correlation between REP and NDVI was high, but decreased as N rate decreased. The decrease in
32
33 correlation could be explained by the reduction of plant biomass which ultimately increases the area of
34
35 soil exposed, and directly influences REP and NDVI values. Some research reports that there is less
36
37 influence of soil background on REP (Meer and Jong, 2006). Therefore, NDVI might be more influenced
38
39 by soil background and as such would behave differently than REP. Detection of biomass amounts using
40
41 different soil backgrounds will require further study.
42
43

44
45 The advantages and/or disadvantages of REP in field based research need to be discussed. Studies
46
47 showed that REP was highly correlated with chlorophyll content at the plant canopy level (Chappelle et
48
49 all.1991; Cho and Skidmore 2006). It was also shown in our research that REP was highly correlated with
50
51 SPAD chlorophyll meter readings. The position and shape of the first derivative spectrum provides more
52
53 opportunity to differentiate plant N response (Cheng et al., 2005; Kupfer and et all., 1990). As illustrated
54
55 in Figure 6, the shape of the first derivative reflectance is more clearly defined in low N rates than high N
56
57 rates. Two points where maximums occur are shown for the 0 kg N/ha rate. At the same plant growth
58
59
60

1
2
3 stage, you can differentiate plant N status by “the shape” of the first derivative reflectance, and can also
4 distinguish “the position” of the maximum point of the first derivative reflectance. The basic point is that
5
6
7 you can manipulate two outcomes, “the position and the shape” of the first derivative reflectance, from
8
9
10 red-edge bands which does not happen with NDVI or simple ratio. Filella and Penuelas (1994) showed
11
12 that the area of the first derivative reflectance has strong correlation with plant biomass. Another
13
14 advantage is that under high biomass, red-edge position could give more accurate estimates of biomass
15
16 (Filella and Penuelas, 1994; Mutanga and Skidmore, 2004). The results show that NDVI is sensitive to
17
18 different plant N response as well as different plant growth stages but the sensitivity tended to be higher
19
20 for REP, especially with advancing growth stage.
21

22
23 There are also disadvantages when using REP. Using the Ocean Optics USB4000 spectrometer,
24
25 REP was very sensitive to noise, or in other words, it is hard to find the position of red edge. Red-edge
26
27 could be obtained using a hyperspectrometer but because of high contents of information per pixel, the
28
29 analysis of derivative system requires the right techniques and time (Ruffin and King, 1999). In this wheat
30
31 study, the range of red-edge position was narrow, not more than 15 nm, between non-N treated plants and
32
33 high N treated plants (Figures 2 and 3). Also as described in Figure 7, some plot samples show that red-
34
35 edge position using the curve fitting method has equal probability for any value wavelength within 710 nm
36
37 to 730nm. To facilitate more simplified capture of information from clearly very narrow wavebands,
38
39 better spectrometers will be required.
40
41

42 CONCLUSIONS

43
44 The REP behaved very similar to NDVI in winter wheat. Because of the costs associated with
45
46 capturing and using REP data in sensor based technology, and since there were no clear benefits over that
47
48 of NDVI, widespread use of REP at present stifled. More evidence from work on a range of crops and is
49
50 needed to verify if REP is significantly better than NDVI. Modifying existing sensors to include REP
51
52 may not be worth the investment in winter wheat. Further research is needed to evaluate REP with
53
54 biomass at early growth stages. In general, these results showed that $NDVI_{RED}$ better detected differences
55
56 in plant growth at the same N rates, especially at early growth stages when compared to REP.
57
58
59
60

REFERENCES


1. Babar, M.A., M. P. Reynolds, M. van Ginkel, A. R. Klatt, W. R. Raun, and M. L. Stone. 2006. Spectral reflectance to estimate genetic variation for in-season biomass, leaf chlorophyll and canopy temperature in wheat. *Crop Science* 46:1046-1057.
2. Bellairs, S.M., N.C. Turner, P.T. Hick, and R.C.G. Smith. 1996. Plant and soil influences on estimating biomass of wheat in plant breeding plots using field spectral radiometers. *Australian Journal of Agricultural Research* 47:1017-1034.
3. Boman., R.K., R.L. Westerman, W.R. Raun. And M.E. Jojola. 1995. Time of nitrogen application: Effects on winter wheat and residual soil nitrate. *Soil Science Society of America Journal* 59:1364-1369.
4. Chappelle, E.W., M.S. Kim, and J.E. McMurtrey III. 1991. The red edge shift: An explanation of its relationship to stress and the concentration of chlorophyll a. Proceedings of the 11th Annual International Geoscience and Remote Sensing Symposium, Espoo, Finland; UNITED STATES; 3-6 June 1991. 2287-2290.
5. Chen, Z., and Elvidge, C. D. 1993. Description of derivative-based high spectral-resolution (AVIRIS) green vegetation index. *Proceedings SPIE* 1937:43-54.
6. Cheng., Y., C. Hu, H. Dai and Y. Lei. 2005. Spectral red edge parameters for winter wheat under different nitrogen support levels. *Proceeding of SPIE-The international society for optical engineering* 5884:384-389.
7. Cho, M.A. and A.K. Skidmore. 2006. A new technique for extracting the red edge position from hyperspectral data: The linear extrapolation method. *Remote Sensing of Environment* 101:181-193.
8. Clevers, J. G. P. W. 1994. Imaging Spectrometry in Agriculture – Plant Vitality and Yield Indicators. *Eurocourses: Remote Sensing* 4:193-219.

- 1
2
3 9. Clevers, J. G. P. W., S. M. de Jong, G. F. Epema, F.V. der Meer, W. H. Bakker, A. K.
4
5 Skidmore and E.A. Addink. 2001. MERIS and the red-edge position. *International Journal of*
6
7 *Applied Earth Observation and Geoinformation* 3:313-320.
8
9
- 10 10. Dawson, T.P., and P.J. Curran. 1998. A new technique for interpolating the reflectance red edge
11
12 position. *International Journal of Remote Sensing* 19:2133-2139.
13
- 14 11. Filella, I. and J. Penuelas. 1994. The red edge position and shape as indicators of plant chlorophyll
15
16 content, biomass and hydric status. *International Journal of Remote Sensing* 15:1459-1470.
17
- 18 12. Fox R.H., W.P. Piekielek, and K.M. Macneal. 1994. Using a chlorophyll meter to predict nitrogen
19
20 fertilizer needs of winter wheat. *Communications Soil Science Plant Analysis* 25:171-181.
21
- 22 13. Gitelson. A.A., Y.J. Kaufman, R. Stark and D.C. Rundquist. 2002. Novel algorithms for remote
23
24 estimation of vegetation fraction. *Remote Sensing of Environment* 80:76–87.
25
- 26 14. Guyot, G., & Baret, F. (1988). Utilisation de la haute résolution spectrale pour suivre l'état des
27
28 couverts végétaux. Proceedings of the 4th International colloquium on spectral signatures of
29
30 objects in remote sensing. ESA SP-287, Assois, France. 279–286.
31
- 32 15. Holtz, S., K. Grima, B. Tubana, D.B. Arnall, D. Edmond, Y. Kanke, J.B. Solie, and W.R. Raun.
33
34 2008. Above ground nitrogen accumulation as a function of time in corn and winter wheat.
35
36 *Journal of Plant Nutrition* (in press).
37
- 38 16. Large, E.C. 1954. Growth stages in cereals: Illustration of the Feekes Scale. *Plant Pathology*
39
40 3:128–129.
41
- 42 17. Kupfer, F.B.G., K. Dockter and W. Kuhbauch. 1990. Shape of the red edge as vitality indicator for
43
44 plants. *International Journal of Remote Sensing* 11:1741-1753.
45
- 46 18. Meer, F.V.D., and S.M. de Jong. 2006. Imaging spectrometry for agriculture applications. In:
47
48 *Imaging spectrometry: Basic principal and prospective application*, eds. Clevers, J. G. P. W. and
49
50 R. Jongschaap, pp.157-197. Dordrecht, Netherlands: Springer.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
19. Mutanga, O. and Skidmore, A.K. 2004. Hyperspectral band depth analysis for a better estimation of grass biomass (*Cenchrus ciliaris*) measured under controlled laboratory conditions. *International Journal of Applied earth Observation and Geoinformation* 5:87-96.
 20. Mutanga, O. and Skidmore, A.K. 2004. Narrow band vegetation indices overcome the saturation problem in biomass estimation. *International Journal of Remote Sensing* 25:3999-4014.
 21. Mutanga, O. and Skidmore, A.K. 2007. Red edge shift and biochemical content in grass canopies. *ISPRS Journal of Photogrammetry and Remote Sensing* 62:34-42.
 22. Raun, W.R., and G.V. Johnson. 1999. Improving nitrogen use efficiency for cereal production. *Agronomy Journal* 91:357-363.
 23. Raun, W.R., J.B. Solie, G.V. Johnson, M.L. Stone, R.W. Mullen, K.W. Freeman, W.E. Thomason, and E.V. Lukina. 2002. Improving nitrogen use efficiency in cereal grain production with optical sensing and variable rate application. *Agronomy Journal* 94:815-820.
 24. Ruffin, C., and King, R. L. 1999. The analysis of hyperspectral data using Savitzky-Golay filtering-Theoretical basis In: IEEE International Geoscience and Remote Sensing Symposium, 1999. CD ROM-IEEE Catalog Number 99CH36293.
 25. Serrano, L., I. Filella, and J. Peñuelas. 2000. Remote sensing of biomass and yield of winter wheat under different nitrogen supplies. *Crop Science* 40:723-731.
 26. Shafri, H.Z.M., M.A.M Salleh and A. Ghiyamat. 2006. Hyperspectral remote sensing of vegetation using red edge position techniques. *American Journal of Applied Sciences* 6:1864-1871.
 27. Shafri, H.Z.M., and M.R.M. Yusof. 2009. Trends and issues in noise reduction for hyperspectral vegetation reflectance spectra. *European Journal of Scientific Research*. 29:404-410.
 28. Solie, J.B., W.R. Raun, R.W. Whitney, M.L. Stone and J.D. Ringer. 1996. Optical sensor based field element size and sensing strategy for nitrogen application. *Transactions of ASABE* 39:1983-1992.
 29. Waldren, R.P and A.D. Flowerday. 1979. Growth stages and distribution of dry matter, N, P, and K in winter wheat. *Agronomy Journal* 71:391-397

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Red-edge position determined by the linear method (Meer and Jong, 2006).



ew Only



Figure 2. NDVI_{RED} and REP plotted against N rate, Stillwater, OK, 2008-2009.


Figure 3. NDVI_{RED} and REP plotted against N rate, Perkins, OK, 2008-2009.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Figure 4. Visual image of winter wheat at Feekes 4 under conventional tillage and no-tillage, Stillwater and Perkins, OK, 2008.



View Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Figure 5. Linear relationship between REP and NDVI_{RED} at Feekes 4 at different N rates, Stillwater, OK, 2008.






Figure 6. Shape of red-edge and its position at Feekes 4 in the 0 kg N/ha and 135 kg N/ha plots, Stillwater, OK, 2008.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 7. Shape of red-edge by derivative method with curve fitting techniques at Feekes 5 in 135 kg N/ha plots, Stillwater, OK, 2009.



view Only

Table 1. Simple correlation coefficients between REP and each index (linear method), Stillwater, OK, 2008-2009.

		NDVI _{RED}	NDVI _{GREEN}	SR _{RED}	SR _{GREEN}	SPAD
2008	Feekes 4	0.766	0.667	0.769	0.677	0.719
	Feekes 5	0.865	0.872	0.914	0.904	0.83
	Feekes 7	0.835	0.918	0.872	0.925	0.846
	Feekes 10	-	-	-	-	-
2009	Feekes 4	0.83	0.767	0.81	0.75	0.326
	Feekes 5	0.774	0.697	0.776	0.697	0.518
	Feekes 7	0.531	0.579	0.48	0.48	0.608
	Feekes 10	0.83	0.814	0.843	0.824	0.637

*Significant at the 0.001 probability level $NDVI_{RED} = \frac{\rho_{780} - \rho_{670}}{\rho_{780} + \rho_{670}}$, $NDVI_{GREEN} = \frac{\rho_{780} - \rho_{560}}{\rho_{780} + \rho_{560}}$, $SR_{RED} = \frac{\rho_{780}}{\rho_{670}}$,

$SR_{GREEN} = \frac{\rho_{780}}{\rho_{560}}$ SPAD=SPAD meter readings.

Table 2. Simple correlation coefficients between NDVI_{RED} and each index, Stillwater, OK, 2008-2009.

		NDVI _{GREEN}	SR _{RED}	SR _{GREEN}	SPAD
2008	Feekes 4	0.978	0.988	0.976	0.482
	Feekes 5	0.978	0.97	0.968	0.787
	Feekes 7	0.956	0.956	0.918	0.709
	Feekes 10	-	-	-	-
2009	Feekes 4	0.956	0.994	0.949	0.396
	Feekes 5	0.986	0.98	0.974	0.367
	Feekes 7	0.962	0.924	0.937	0.345
	Feekes 10	0.988	0.966	0.97	0.408

*Significant at the 0.001 probability except SPAD, $NDVI_{GREEN} = \frac{\rho_{780} - \rho_{560}}{\rho_{780} + \rho_{560}}$, $SR_{RED} = \frac{\rho_{780}}{\rho_{670}}$, $SR_{GREEN} = \frac{\rho_{780}}{\rho_{560}}$

SPAD=SPAD meter readings.

Table 3. Simple correlation coefficients between REP and each index (linear method), Perkins, OK, 2008-2009.

		NDVI _{RED}	NDVI _{GREEN}	SR _{RED}	SR _{GREEN}	SPAD
2008	Feekes 4	0.924	0.918	0.887	0.878	0.238
	Feekes 5	0.863	0.878	0.83	0.776	0.265
	Feekes 7	-	-	-	-	-
	Feekes 10	0.941	0.953	0.931	0.924	0.261
2009	Feekes 4	0.686	0.593	0.618	0.702	0.74
	Feekes 5	-	-	-	-	-
	Feekes 7	0.955	0.955	0.941	0.935	0.745
	Feekes 10	0.814	0.856	0.771	0.748	0.812

*Significant at the 0.001 probability level except SPAD, $NDVI_{RED} = \frac{\rho_{780} - \rho_{670}}{\rho_{780} + \rho_{670}}$, $NDVI_{GREEN} = \frac{\rho_{780} - \rho_{560}}{\rho_{780} + \rho_{560}}$,

$SR_{RED} = \frac{\rho_{780}}{\rho_{670}}$, $SR_{GREEN} = \frac{\rho_{780}}{\rho_{560}}$ SPAD=SPAD meter readings.

Table 4. Simple correlation coefficients between NDVI_{RED} and each index, Perkins, OK, 2008-2009.

		NDVI _{GREEN}	SR _{RED}	SR _{GREEN}	SPAD
2008	Feekes 4	0.994	0.99	0.99	0.149
	Feekes 5	0.994	0.974	0.951	0.108
	Feekes 7	-	-	-	-
	Feekes 10	0.99	0.962	0.968	0.199
2009	Feekes 4	0.982	0.978	0.992	0.918
	Feekes 5	-	-	-	-
	Feekes 7	0.994	0.972	0.972	0.689
	Feekes 10	0.978	0.893	0.92	0.75

*Significant at the 0.001 probability except SPAD, $NDVI_{GREEN} = \frac{\rho_{780} - \rho_{560}}{\rho_{780} + \rho_{560}}$, $SR_{RED} = \frac{\rho_{780}}{\rho_{670}}$, $SR_{GREEN} = \frac{\rho_{780}}{\rho_{560}}$

SPAD=SPAD meter readings.

Table 5. Model parameters between REP and different N rates or between NDVI_{RED} and different N rates.

	Variable	Index	Mean	Std Dev	Std Error
Linear (y=ax+b)	r ²	NDVI _{RED}	0.661	0.201	0.058
	r ²	REP	0.733	0.112	0.032
	r ²	Difference (NDVI _{RED} -REP)	-0.071	0.163	0.067
	Adjusted r ²	NDVI _{RED}	0.6125	0.2619	0.0726
	Adjusted r ²	REP	0.6632	0.1684	0.0467
	Adjusted r ²	Difference (NDVI _{RED} -REP)	-0.0507	0.2202	0.0864
	Standard Error	NDVI _{RED}	0.0545	0.0203	0.00564
	Standard Error	REP	1.1899	0.5836	0.1619
	Standard Error	Difference (NDVI _{RED} -REP)	-1.1354	0.4129	0.162
	F score	NDVI _{RED}	59.1929	44.1177	12.2361
	F score	REP	72.5746	66.0809	18.3275
	F score	Difference (NDVI _{RED} -REP)	-13.3817	56.183	22.0368
Logarithm (y-ax+b ln(x))	r ²	NDVI _{RED}	0.6422	0.2126	0.059
	r ²	REP	0.7077	0.1338	0.0371
	r ²	Difference (NDVI _{RED} -REP)	-0.0656	0.1776	0.0697
	Adjusted r ²	NDVI _{RED}	0.5926	0.2513	0.0697
	Adjusted r ²	REP	0.6625	0.1661	0.0461
	Adjusted r ²	Difference (NDVI _{RED} -REP)	-0.0699	0.213	0.0835
	Standard Error	NDVI _{RED}	0.0564	0.02	0.00555
	Standard Error	REP	1.1783	0.5585	0.1549
	Standard Error	Difference (NDVI _{RED} -REP)	-1.1219	0.3952	0.155
	F score	NDVI _{RED}	52.9866	42.0479	11.662
	F score	REP	77.7351	86.0683	23.871
	F score	Difference (NDVI _{RED} -REP)	-24.7485	67.7339	26.5674
Exponential (y=a*exp(bx))	r ²	NDVI _{RED}	0.6614	0.2011	0.0581
	r ²	REP	0.7328	0.1128	0.0326
	r ²	Difference (NDVI _{RED} -REP)	-0.0714	0.1631	0.0666
	Adjusted r ²	NDVI _{RED}	0.6179	0.2336	0.0674
	Adjusted r ²	REP	0.694	0.1399	0.0404
	Adjusted r ²	Difference (NDVI _{RED} -REP)	-0.0761	0.1926	0.0786
	Standard Error	NDVI _{RED}	0.0571	0.0215	0.00621
	Standard Error	REP	1.1566	0.594	0.1715
	Standard Error	Difference (NDVI _{RED} -REP)	-1.0995	0.4203	0.1716
	F score	NDVI _{RED}	55.7023	42.5734	12.2899

	F score	REP	78.5833	65.9384	19.0348
	F score	Difference (NDVI_{RED}-REP)	-22.881	55.4994	22.6575
Sigmoid	r²	NDVI_{RED}	0.725	0.2133	0.0592
	r²	REP	0.7732	0.1515	0.0392
	r²	Difference (NDVI_{RED}-REP)	-0.048	0.181	0.071
	Adjusted r²	NDVI_{RED}	0.6467	0.2799	0.0776
	Adjusted r²	REP	0.6866	0.228	0.632
	Adjusted r²	Difference (NDVI_{RED}-REP)	-0.04	0.2553	0.1001
	Standard Error	NDVI_{RED}	0.0503	0.0233	0.0065
	Standard Error	REP	1.0705	0.4432	0.1229
	Standard Error	Difference (NDVI_{RED}-REP)	-1.02	0.3138	0.1231
	F score	NDVI_{RED}	29.564	26.349	7.3079
	F score	REP	33.732	35.379	9.8124
	F score	Difference (NDVI_{RED}-REP)	-4.168	31.192	12.235

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Table 6. Results of T-test between REP and NDVI models.

	Parameter	df	t value	P> t
Linear (y=ax+b)	r²	24	-0.68	0.09
	Adjusted r²	24	-0.59	0.56
	Standard Error	24	-7.01	<0.0001
	F score	24	-0.61	0.55
Logarithm (y-ax+b ln(x))	r²	24	-0.94	0.36
	Adjusted r²	24	-0.84	0.41
	Standard Error	24	-7.24	<0.0001
	F score	24	-0.93	0.36
Exponential (y=a*exp(bx))	r²	22	-1.07	0.3
	Adjusted r²	22	-0.97	0.34
	Standard Error	22	-6.41	<0.0001
	F score	22	-1.01	0.32
Sigmoid	r²	21	0.75	0.46
	Adjusted r²	21	1.15	0.03
	Standard Error	21	-6.93	<0.0001
	F score	21	0.1	0.92



AGREEMENT IN RELATION TO COPYRIGHT IN AN ARTICLE FOR A TAYLOR & FRANCIS/ROUTLEDGE JOURNAL

In order to ensure both the widest dissemination and protection of material published in our Journal, we ask Authors to transfer to the Publisher, Taylor & Francis, the rights of copyright in the Articles they contribute. This enables Taylor & Francis to ensure protection against infringement. The transfer of copyright must be clearly stated in writing.

PLEASE PROVIDE US WITH THE FOLLOWING INFORMATION, REVIEW OUR POLICIES, AND CONFIRM YOUR ACCEPTANCE OF THE TERMS OF THE ATTACHED ARTICLE PUBLISHING AGREEMENT BY SIGNING THIS FORM AS INDICATED BELOW.

Article (the "Article") entitled: Red Edge as a Potential Index for Detecting Differences in Plant Nitrogen Status in Winter Wheat
Author(s): Kanke, Yumiko, William Raus, John Solie, Marvin Stone, Randal Taylor
To be published in the journal (the "Journal"): Journal of Plant Nutrition [ISSN: 0190-4167]

YOUR STATUS

- I am the sole author of the Article
Please indicate if any of the statements below also apply to you:
I am a UK, Canadian or Australian Government employee and claim Crown Copyright
I am a US Government employee and there is no copyright to transfer
I am an NIH employee and there is no copyright to transfer. I submit this form together with an NIH addendum.
I am a contractor of the US Government (includes NIH contractors) under contract number:
I am required to sign this form
I am one of multiple co-authors of the Article and confirm I have the consent of my co-authors to sign this agreement on their behalf
Please indicate if any of the below also apply to you and your co-authors:
All of my co-authors are UK, Canadian or Australian Government employees and Crown Copyright is claimed / not claimed (circle one)
One or more of my co-authors, but not all of them, are UK, Canadian or Australian Government employees and Crown Copyright is claimed / not claimed (circle one)
All of my co-authors are US Governmental employees and there is no copyright to transfer
The work was performed by contractors of the US Government under contract number
The copyright in the Article belongs to my employer (is a "work made for hire") and I am granting licence to publish as an authorized representative of my employer. My Title and Company are stated in the section below.

ASSIGNMENT OF PUBLISHING RIGHTS

I hereby assign to Taylor & Francis the copyright in the above specified manuscript (government authors not transferring copyright hereby assign a non-exclusive licence to publish) and any accompanying tables, illustrations, data and any other supplementary information intended for publication in all forms and all media (whether known at this time or developed at any time in the future) throughout the world, in all languages, for the full term of copyright, to take effect if and when the article is accepted for publication. If I am one of several co-authors, I hereby confirm that I am authorized by my co-authors to grant this Licence as their agent on their behalf. For the avoidance of doubt, this assignment includes the rights to supply the article in electronic and online forms and systems.

I confirm that I have read and accept the full terms of the Journal's article publishing agreement attached to this form including my author warranties, and have reviewed the Journal's policies on Author Rights.

Signed: William Raus Name Printed: William Raus
Title and Company (if employer representative): Regents Professor Oklahoma State University Date: May 24, 2011

Please return only this page completed and physically signed. You may submit by fax, postal mail, email, or upload to CATS.

THIS FORM WILL BE RETAINED BY THE PUBLISHER.